Problem K Continued Fraction Problem ID: fraction Time Limit: 1 second

The (simple) continued fraction representation of a real number r is an expression obtained by an iterative process of representing r as the sum of its integer part and the reciprocal of another number, then writing this other number as the sum of its integer part and another reciprocal, and so on. In other words, a continued fraction representation of r is of the form

$$r = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}$$



where a_0, a_1, a_2, \ldots are integers and $a_1, a_2, \cdots > 0$. We call the a_i -

values *partial quotients*. For example, in the continued fraction representation of 5.4, the partial quotients are $a_0 = 5$, $a_1 = 2$, $a_2 = 2$. This representation of a real number has several applications in theory and practice. If r is a rational number, the partial quotients are eventually all zero, so we only need to consider a finite number of partial quotients.

Given two rational numbers in continued fraction representation, your task is to perform the four elementary arithmetic operations on these numbers and display the results in continued fraction representation.

Input

The input consists of a single test case. The test case consists of three lines. The first line contains two integers n_1 and n_2 , where $1 \le n_i \le 9$ is the number of partial quotients of rational number r_i for $1 \le i \le 2$. The second line contains the partial quotients of r_1 and the third line contains the partial quotients of r_2 . The absolute values of the quotients are not more than 10 and you may assume that $r_1 > r_2 > 0$.

Output

Display the partial quotients of the continued fraction representations of $r_1 + r_2$, $r_1 - r_2$, $r_1 \times r_2$, and r_1/r_2 , in order, each in a line. Consecutive partial quotients on each line are separated by a single space. Do not print any trailing zero partial quotients.

Sample Input	Sample Output
4 3	11
5 1 1 2	0 5
5 2 2	30 4 6
	1 27